Sports Psychiatry: A Mental Health Primer for College and Professional Athletes

Stacy Leon, MD[1]* and John Powers[1]*

[1]Co–First Authors

*Stacy Leon, MD and John Powers contributed equally as first authors.

This guide is intended for athletes, coaches, teams, and clinicians interested in optimizing mental health and performance through the lens of sports psychiatry. All rights reserved. For educational use only.

Introduction

Elite athletes are often perceived as the epitome of mental toughness and resilience. But behind the highlight reels and medals lie immense pressures; expectations to perform, intense training schedules, media scrutiny, injuries, and the often-overlooked emotional toll of transitions in and out of sport. These factors contribute to real and significant mental health challenges.

This guide on sports psychiatry, performance, and mental wellness provides essential information for college and professional athletes, along with the coaches, clinicians, and support teams around them..

Mental health symptoms in athletes are often hidden or misunderstood. Elite sports require enormous time, effort, and financial commitment to reach the top. While most athletes say they're willing to "do anything" to succeed, few are fully aware of the risks those sacrifices pose to both mental and physical health. Left unchecked, these pressures can lead to burnout, injury, or a serious drop in performance. That's why it's crucial that athletes and the biopsychosocial systems they engage with understand how these stressors affect both health and performance.

Understanding Mental Health in Athletes

Structural Influences on Mental Health

Athletes don't just train their bodies. They are shaped by the environments and systems around them. One of the most influential of these is coaching culture. While coaches often aim to instill mental toughness, discipline, and resilience, these traits are sometimes enforced through outdated methods: verbal abuse, humiliation during practice, and punishment-based training.

While the goal may be to build strength, these techniques can create lasting harm, on the court and beyond. Athletes often carry the emotional residue from training into their home and social lives. Coaches frequently pass down the same methods they experienced as players, believing "it worked for me, so it'll work for them," without recognizing the long-term mental health consequences.

Prevalent Disorders in Elite Athletes

Anxiety Disorders

Anxiety is the most commonly reported mental health condition among elite athletes across sports and levels of competition. It may present as generalized anxiety, performance anxiety, panic attacks, or social anxiety, and can have a direct impact on focus, confidence, and physical performance.

Research from the International Olympic Committee (IOC) emphasizes that anxiety symptoms are highly prevalent in athletes, particularly in the lead-up to major competitions and during transitions such as injury, team changes, or retirement (Reardon et al., 2019).

According to the NCAA Mental Health Best Practices report, student-athletes face unique stressors including balancing academics and sport, maintaining scholarships, and performing under public scrutiny; all of which increase the risk for anxiety and related disorders (NCAA, 2021).

Anxiety appears to be reported more frequently by female athletes, yet this may reflect underreporting by male athletes due to cultural norms that discourage emotional expression in men (Rice et al., 2016). As the DSM-5 outlines, symptoms of anxiety can be easily masked by irritability, fatigue, or physical complaints, which are often misattributed in athletic settings (American Psychiatric Association, 2013).

The IOC SMHAT-1 tool was developed to help detect early signs of anxiety and other mental health concerns through structured, sport-specific screening emphasizing early detection and integrated care (Gouttebarge et al., 2021).

Eating Disorders

Eating disorders are a significant concern in elite sports, with prevalence rates notably higher among athletes than in the general population. Research suggests that up to 45% of female athletes and 19% of male athletes report symptoms of disordered eating or clinical eating disorders (The Emily Program, 2023; National Eating Disorders Association, 2021).

The risk is particularly elevated in sports where weight, physique, or aesthetics are emphasized, including:

- **Aesthetic sports** (e.g., gymnastics, figure skating, ballet): Emphasis on thinness and visual appearance contributes to disordered eating behaviors.
- **Endurance sports** (e.g., long-distance running, swimming): High energy demands combined with body weight pressures increase susceptibility.
- **Weight-class sports** (e.g., wrestling, boxing): Athletes often engage in extreme weight-cutting practices to meet strict competition requirements (National Eating Disorders Association, 2021).

Common eating disorders in athletic populations include:

- **Anorexia nervosa**: Severe caloric restriction paired with an intense fear of weight gain.
- **Bulimia nervosa**: Cycles of binge eating followed by compensatory behaviors such as vomiting or overexercising.
- **Anorexia athletica**: A non-clinical term describing compulsive exercise and restrictive eating habits aimed at enhancing performance (The Emily Program, 2023).

These disorders can lead to significant physical and psychological harm, including hormonal imbalances, decreased bone density, cardiovascular issues, and increased injury risk. They can also impair athletic performance by reducing energy availability, focus, endurance, and overall recovery (National Eating Disorders Association, 2021).

Substance Use Disorders

Substance misuse among elite athletes can arise from many factors, whether it's to gain a competitive edge, manage pain, or cope with anxiety, stress, or emotional fatigue. The substances used can range from performance-enhancing drugs (like anabolic steroids or stimulants) to opioids, alcohol, and cannabis.

Some athletes begin using substances with a clear performance goal in mind, while others start with medically prescribed medications that evolve into dependence. The pressure to “play through pain” or constantly meet expectations can normalize misuse within locker rooms or among teammates.

Periods of transition, such as recovering from injury, being cut from a team, or retiring from sport, can also increase vulnerability. During these times, athletes may feel isolated or uncertain about their identity, making them more susceptible to unhealthy coping mechanisms.

Research shows:

- Athletes use performance-enhancing drugs for strength, endurance, and faster recovery, but many of these substances carry physical and psychological risks (Petroczi & Naughton, 2008).
- The risk of opioid misuse rises when athletes are prescribed painkillers for sports injuries, especially without mental health support (Gouttebarge et al., 2021).
- A 2022 report found that over 500 professional football players in the UK sought help for addiction related to drugs, alcohol, and gambling (The Sun, 2022).

To address these risks, early intervention and mental health support are critical. Education about the consequences of substance misuse, confidential access to care, and a culture that values mental wellness over performance alone can make a meaningful difference.

Attention-Deficit/Hyperactivity Disorder (ADHD)

Sports can uniquely mask the symptoms of ADHD due to the natural positive reinforcement provided by physical activity. Many individuals with ADHD are drawn to high-intensity or fast-paced sports such as basketball, football, or tennis, which demand quick decision-making and constant stimulation. These environments can temporarily improve focus, providing a form of behavioral regulation through movement and reward-based feedback (Reardon et al., 2019).

However, while athletic performance may appear unaffected on the surface, athletes with ADHD often struggle to manage other aspects of life, such as academics, relationships, or emotional regulation. These challenges are especially pronounced in college athletes who must balance demanding training schedules with academic performance (NCAA, 2021).

ADHD treatment in elite sports presents unique challenges. Common pharmacological treatments, such as amphetamine-based stimulants, are often classified as performance-enhancing substances by athletic governing bodies. This classification can create a barrier to care, leaving athletes to navigate a difficult balance between managing their symptoms and staying compliant with anti-doping regulations (Gouttebarge et al., 2021).

Despite these concerns, therapeutic enhancement of focus and self-regulation is not the same as ergogenic performance enhancement. The IOC consensus highlights that properly prescribed and monitored treatment—including medication, cognitive behavioral therapy (CBT), and psychoeducation can improve quality of life without conferring unfair athletic advantages (Reardon et al., 2019; Rice et al., 2016).

CBT in particular has shown benefit in helping athletes with ADHD develop coping strategies, improve focus, and manage emotional reactivity, especially when used in combination with support from sports psychiatrists or psychologists (American Psychiatric Association, 2013).

Depression

Depression is one of the most under-recognized and underreported mental health conditions in elite athletes. While athletes are often perceived as highly motivated, goal-driven individuals, the internal experience of depression can be masked by outward success, discipline, or fatigue from training. Symptoms like persistent low mood, loss of interest, poor sleep, or changes in appetite are frequently misattributed to overtraining, burnout, or poor performance (American Psychiatric Association, 2013).

Studies suggest that elite athletes experience depression at rates similar to or greater than the general population. The IOC consensus statement notes that up to 34% of current elite athletes may experience symptoms of depression at some point in their careers (Reardon et al., 2019). Depression is also more commonly reported in female athletes than in males—though underreporting by male athletes due to stigma remains a concern (Rice et al., 2016).
Risk factors include:

- Injury and prolonged recovery periods
- Loss of athletic identity during transitions or retirement
- Performance slumps and fear of failure
- External pressures from media, coaches, or sponsors
- Academic stress in student-athletes (NCAA, 2021)

Despite these challenges, early recognition and treatment of depression can significantly improve both quality of life and performance outcomes. Effective approaches include psychotherapy (such as cognitive behavioral therapy or psychodynamic therapy), medication when appropriate, and strong social support from teammates, coaches, family, and mental health professionals (Gouttebarge et al., 2021; Reardon et al., 2019).

Promoting a team culture that values vulnerability and mental health is key. Depression is treatable, and addressing it early can protect not only an athlete's mental well-being but also their longevity in sport.

Unique Stressors for College and Professional Athletes

College and professional athletes face a range of stressors that extend beyond the playing field. Many must balance the demands of sport with academic responsibilities or evolving career goals. The constant threat of injury, or the reality of physical trauma can create ongoing anxiety; while transitions between teams, roles, or career stages often bring instability. In today's digital age, athletes are also subject to intense public scrutiny, media attention, and the pressures of maintaining a presence on social media. Financial concerns, especially during periods of transition or uncertain contracts, add yet another layer of stress. These challenges often stack on top of one another like layers, compounding over time and making them increasingly difficult to manage. Understanding these stressors, and learning how to work through them, is essential to protecting long-term mental health and optimizing performance.

Supporting Athlete Mental Health

Recognizing Signs and Symptoms

Mental health struggles in athletes often go unnoticed, hidden behind discipline, performance, or the expectation to "push through." But recognizing early warning signs is key to timely support and recovery.

Psychological signs may include mood swings, irritability, persistent sadness or anxiety, withdrawal from teammates, or a noticeable drop in motivation and focus. Athletes may also lose enjoyment in their sport or struggle to make decisions.

Physical and behavioral signs can involve changes in appetite or weight, sleep problems like insomnia or oversleeping, substance misuse, or even self-harm behaviors. Underperformance on the field is often one of the earliest signs.

As emphasized by the IOC and NCAA, recognizing these symptoms, whether subtle or severe, can help athletes get the support they need before issues worsen (Reardon et al., 2019; NCAA, 2021).

Support and Resources

Building a strong support system is essential for athlete mental health. A well-rounded mental health team helps address both performance challenges and deeper emotional concerns.

Sports Psychiatrist – A medical doctor who can diagnose and treat mental illness, prescribe medication, and support performance, behavioral, and cognitive concerns. Sports psychiatrists are uniquely positioned to care for the whole athlete, from mindset to mental health.

Sports Psychologist – Specializes in performance, motivation, stress management, and behavioral strategies. Often works alongside psychiatrists and coaches to optimize mental resilience.

Athletic Trainers and Coaches – Frequently the first to notice changes in mood, behavior, or performance. Their role in early recognition and referral is vital.

Teammates and Family – Trusted relationships can offer day-to-day emotional support and are often key to encouraging athletes to seek help.

Seeking Help

Asking for help is not a sign of weakness, it's a sign of awareness and strength. In high-performance environments, athletes are often conditioned to push through pain, but mental health deserves the same attention and care as physical injuries.

To support help-seeking:

- **Normalize mental health conversations** in team settings, just like injury prevention or nutrition.
- **Make resources visible and accessible**, including contact information for mental health professionals.
- **Offer confidential pathways** for athletes to seek care without fear of judgment or career consequences.
- **Encourage regular mental health check-ins**, even when things seem fine—early support can prevent bigger issues later on.

The sooner support is accessed, the better the outcome for both well-being and performance.

Treatment Approaches

Psychotherapy

Evidence-based therapies like **Cognitive Behavioral Therapy (CBT)** can improve both mood and performance. **Acceptance and Commitment Therapy (ACT)** is useful for navigating identity issues and life transitions, while **psychodynamic therapy** can help athletes explore deeper stressors related to relationships, self-worth, or the demands of elite sport. For those with trauma histories, **trauma-informed care** ensures treatment is both safe and sensitive.

Medication

Medications such as **SSRIs or SNRIs** are commonly used to treat anxiety and depression. **Stimulant and non-stimulant options** are available for managing ADHD. Anti-anxiety medications may be used selectively, with careful attention to side effects and potential impact on performance.

Integrated Care

The best outcomes often come from a **team-based approach**, involving psychiatrists, psychologists, nutritionists, and medical providers. Ongoing communication helps monitor how treatments, especially medications, affect sleep, focus, and physical performance.

Mental Health Screening Tools

Screening tools can help detect early signs of mental health challenges and guide referrals. These are not diagnostic but can prompt further evaluation by professionals.

- **PHQ-9** (Patient Health Questionnaire-9) – screens for depression
- **GAD-7** (Generalized Anxiety Disorder-7) – screens for anxiety
- **AUDIT-C** – brief screening for alcohol use
- **Athlete-Specific Measures** – customized tools to assess stress, identity, and burnout

These tools should be administered confidentially and ideally interpreted by trained professionals. For full screening questions, see Appendix.

Athlete Cheat Sheet: Mental Health Quick Guide

Top 5 Mental Health Tips

1. Check in with yourself daily on mood, stress, sleep, energy.
2. Don't ignore mental or emotional pain, early support is key.
3. Sleep is recovery, make it a priority.
4. Normalize seeking help. Your brain is part of your body.
5. Stay connected because relationships are protective.

Emergency Signs

- Thoughts of self-harm or suicide
- Severe depression or panic attacks
- Dangerous substance use

If you or someone you know is in crisis, seek help immediately: call or text 988 in the U.S.

Quick Contacts Template

- Mental Health Provider: _____
- Athletic Trainer: _____
- Campus/Team Support Line: _____
- Emergency Contact: _____

References

American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). https://doi.org/10.1176/appi.books.9780890425596

Gouttebarge, V., Bindra, A., Blauwet, C., Campriani, N., Hainline, B., Hitchcock, M. E., ... & Reardon, C. L. (2021). International Olympic Committee (IOC) Sport Mental Health Assessment Tool 1 (SMHAT-1). *British Journal of Sports Medicine, 55*(1), 30–37. https://doi.org/10.1136/bjsports-2020-102411

NCAA Sport Science Institute. (2021). *Mental health best practices: Inter-association consensus document*. https://www.ncaa.org/sports/2021/1/11/mental-health-best-practices.aspx

National Eating Disorders Association. (2021). *Eating disorders in athletes*. https://www.nationaleatingdisorders.org/eating-disorders-athletes

Petroczi, A., & Naughton, D. P. (2008). Performance enhancement with supplements: Incongruence between rationale and practice. *Journal of the International Society of Sports Nutrition, 5*, 22. https://doi.org/10.1186/1550-2783-5-22

Reardon, C. L., Hainline, B., Aron, C. M., Baron, D., Baum, A. L., Bindra, A., ... & Engebretsen, L. (2019). Mental health in elite athletes: International Olympic Committee consensus statement. *British Journal of Sports Medicine, 53*(11), 667–699. https://doi.org/10.1136/bjsports-2019-100715

Rice, S. M., Purcell, R., De Silva, S., Mawren, D., McGorry, P. D., & Parker, A. G. (2016). The mental health of elite athletes: A narrative systematic review. *Sports Medicine, 46*(9), 1333–1353. https://doi.org/10.1007/s40279-016-0492-2

The Emily Program. (2023). *Eating disorders in athletes: Understanding the risks*. https://www.emilyprogram.com/blog/eating-disorders-in-athletes-understanding-the-risks/

The Sun. (2022, March 10). *Over 500 footballers treated for addiction to alcohol, drugs and gambling*. https://www.thesun.co.uk/sport/football/17915246/footballers-addiction-drugs-alcohol-gambling/

Appendix: Screening Tools

PHQ-9 (Patient Health Questionnaire-9)

Scoring: 0 (Not at all) to 3 (Nearly every day); total score range: 0–27
Over the last 2 weeks, how often have you been bothered by the following problems?

1. Little interest or pleasure in doing things
2. Feeling down, depressed, or hopeless
3. Trouble falling or staying asleep, or sleeping too much
4. Feeling tired or having little energy
5. Poor appetite or overeating
6. Feeling bad about yourself — or that you are a failure or have let yourself or your family down
7. Trouble concentrating on things, such as reading the newspaper or watching television
8. Moving or speaking so slowly that other people could have noticed? Or the opposite — being so fidgety or restless that you have been moving a lot more than usual
9. Thoughts that you would be better off dead or of hurting yourself in some way

GAD-7 (Generalized Anxiety Disorder-7)

Scoring: 0 (Not at all) to 3 (Nearly every day); total score range: 0–21
Over the last 2 weeks, how often have you been bothered by the following problems?

1. Feeling nervous, anxious, or on edge
2. Not being able to stop or control worrying
3. Worrying too much about different things
4. Trouble relaxing
5. Being so restless that it is hard to sit still
6. Becoming easily annoyed or irritable
7. Feeling afraid as if something awful might happen

AUDIT-C (Alcohol Use Disorders Identification Test - Consumption)

Scoring: 0–12; scores $\geq$4 in men or $\geq$3 in women suggest possible alcohol misuse

1. How often do you have a drink containing alcohol?
2. How many standard drinks containing alcohol do you have on a typical day?
3. How often do you have six or more drinks on one occasion?

Athlete-Specific Screening Prompts

- How often do you feel pressure to perform from coaches, teammates, or sponsors?
- Do you ever feel that your identity is too closely tied to your sport?
- Have you experienced burnout or loss of enjoyment in your sport?
- Have you ever hidden an injury or emotional struggle to avoid being benched or judged?

These tools can provide a helpful starting point but should always be followed up with a conversation and professional evaluation.

This primer is a living document and should evolve with the athlete's needs and emerging research in sports psychiatry. By prioritizing mental health, athletes can achieve sustainable performance and long-term wellbeing.